



Biomonitoring exposure to environmental tobacco smoke (ETS): A critical reappraisal

G Scherer¹ and E Richter²

¹Analytisch-biologisches Forschungslabor München (ABF), Goethestrasse 20, D-80336 München; ²Walther-Straub-Institut für Pharmakologie und Toxikologie der Universität München, Nussbaumstrasse 26, D-80336 München, Germany

- 1 The most frequently used biomarkers for exposure to environmental tobacco smoke (ETS) are cotinine and thiocyanate in body fluids, carboxyhaemoglobin in red blood cells (COHb) and carbon monoxide in the expired air. Although not ideal, cotinine in blood, saliva or urine is an established biomarker for ETS exposure within the past 1–3 days. Comparison with cotinine concentrations in cigarette smokers reveals that passive smokers take up less than 1/100 of the nicotine dose of smokers.
- 2 Biomonitoring data available for the ETS-related exposure to genotoxic substances comprise uptake of benzene, polycyclic aromatic hydrocarbons (PAH), aromatic amines, tobacco-specific nitrosamines (TSNA), electrophilic compounds giving rise to urinary thioethers, mutagens causing urinary mutagenic activity and the formation of various DNA adducts. With the exception of TSNA, these biomarkers are related to chemicals occurring ubiquitously in the environment and in the food. As a consequence, the background levels in unexposed nonsmokers are high compared to the observed increases (if any) associated with ETS exposure.
- 3 Some markers of biological effects, which, by definition, are non-specific with regard to the underlying exposure, have also been investigated in relation to ETS exposure. These markers comprise cytogenetic effects, aryl hydrocarbon hydroxylase (AHH) induction, urinary hydroxyproline excretion and various factors indicative of cardiovascular risks. The available data suggest that passive smoking is associated with a small induction of placental AHH and also with effects on cardiovascular risk markers. The latter findings in particular may be confounded by other risk factors, which have been observed to be more frequent in passive smokers than in unexposed nonsmokers.

Keywords: biomonitoring; environmental tobacco smoke; cotinine; thiocyanate; carbon monoxide; benzene

Introduction

Quantitative risk estimates for ETS-exposed nonsmokers published by various authorities rely almost exclusively on data from epidemiological studies.¹ It is the purpose of this review to summarize available data on the internal dose of tobacco smoke constituents and their possible effects in nonsmokers, exposed to ETS under real-life conditions. We feel that biological monitoring with nonsmokers exposed to ETS in their normal environment can provide objective data in this controversial field of research.

Surrogate biomarkers for ETS exposure

Surrogate biomarkers for ETS exposure are markers related to substances in tobacco smoke, which are assumed not to be implicated in toxicologically

relevant processes, e.g. cancer, cardiovascular diseases or respiratory dysfunctions. The concentrations of precursors of surrogate biomarkers in ETS should be directly proportional to those of toxicologically relevant constituents in ETS. Suitable biomarkers should be specific for the exposure, their biological half-life should be long enough and the analytical method should be specific, precise, and not too labourious and expensive.² Strictly, these criteria are not met by any of the most commonly used surrogate biomarkers such as carboxyhaemoglobin or carbon monoxide in the exhalate, as well as cotinine and thiocyanate in body fluids (Table 1).

Carboxyhaemoglobin (COHb) and exhaled CO (COex)

Carbon monoxide (CO) is generated during all incomplete combustions of organic materials (for example cooking, heating or vehicle exhaust). In addition, CO is endogenously formed in mammals during catabolism of haem moieties.³ Therefore, the biomarkers COHb and COex are by no means specific for tobacco smoke exposure. Normal or

Correspondence: G Scherer
Received 3 March 1997; revised 2 April 1997; accepted 4 April 1997

Table 1. Surrogate biomarkers for ETS exposure^a

Biomarker	Precursor in ETS	Other sources	Significant association with the extent of real-life ETS exposure ^b
COHb, COex	CO	Incomplete combustions, endogenous formation	No ^{109,110}
Thiocyanate in body fluids	HCN	Diet	No ^{1,111,112}
Cotinine in body fluids	Nicotine	Diet(?)	Yes ^{12,113,114}

^aAbbreviations: COHb: carboxyhaemoglobin; COex: carbon monoxide in exhaled air. ^bLiterature cited is not comprehensive; for a review, see Benowitz.¹⁶

only marginally increased levels of COHb or COex were observed in passive smokers as compared to nonsmokers. This is compatible with the finding that the ETS-related increase of CO indoors amounts to only 0.5–1 p.p.m.⁴

Thiocyanate in body fluids

Thiocyanate is the detoxification product of cyanide with a half-life of 10–14 days.⁵ The precursor of thiocyanate in tobacco smoke is hydrogen cyanide. Hydrogen cyanide levels in mainstream smoke (MS) of cigarettes vary between 300 and 550 µg/cigarette and the ratio of sidestream smoke (SS) to MS is 0.19–0.37.⁶ In ETS, hydrogen cyanide occurs almost exclusively in the gaseous phase. Other sources for the intake of cyanide are almonds, pulses and maize. Cyanide is also formed in the colon by bacteria. Preformed thiocyanates are found in cabbage, turnips, mustard and cow milk.⁴ Therefore, thiocyanate in body fluids is not specific for exposure to tobacco smoke. Although thiocyanate can be easily measured, and its long biological half-life prevents large fluctuations in body fluids, it is not specific enough to be suitable as a biomarker for ETS exposure under real-life conditions.

Cotinine in body fluids

Cotinine is the main metabolite of nicotine, the principal alkaloid of the tobacco plant. The nicotine yield of cigarette MS is about 1 mg, and the SS/MS ratio amounts to 2.6–3.3.⁶ Nicotine occurs almost completely in the particulate phase of mainstream smoke and in the gaseous phase of ETS.⁷ Average nicotine concentrations in indoor environments where smoking occurs are usually in the range of 1–10 µg/m³.^{6,9} In addition, some *solanaceae*-derived food items such as tomatoes, potatoes and egg plants as well as tea contain small amounts of nicotine.¹⁰ The biological half-life of cotinine is considerably longer than that of nicotine (16–20 h vs 1 h).¹¹ Therefore, cotinine is a much more reliable

biomarker for active smoking, as well as ETS exposure, than is nicotine. In almost all studies, a statistically significant difference in cotinine levels between ETS-exposed and nonexposed nonsmokers was observed (Table 1). In addition, a significant relationship between the cotinine concentration in body fluids and the extent (frequency, duration and intensity) of ETS exposure was found. ETS exposure at home was often found to be more important than ETS exposure at the workplace or other places.^{9,12} The ratio between the cotinine concentrations in ETS-exposed nonsmokers and smokers is usually <1/100. However, there are general limitations for comparisons of active smoking with ETS exposure.¹³ A major difference is the fact that cotinine in smokers is an indicator of the nicotine uptake with the particulate matter of mainstream smoke, whereas cotinine in nonsmokers indicates the exposure to the ETS gaseous phase.¹⁴ The longer biological half-life of cotinine in nonsmokers as compared to smokers would lead to an overestimation of the exposure in passive smokers relative to smokers.¹⁵ However, other researchers found similar pharmacokinetics of nicotine and cotinine in smokers and nonsmokers.¹⁶ On the other hand, the significantly shorter half-life of airborne nicotine compared to other ETS components could lead to an underestimation of ETS exposure when based on cotinine concentrations in body fluids.¹⁷ Airborne nicotine shows a high degree of adsorption to surfaces in indoor environments, which is responsible for the high decay rates of nicotine. Adsorbed nicotine can be released into the air so that nicotine, but not other ETS constituents, are measurable in environments in the absence of smoking.¹⁸ This, together with possible dietary nicotine intake and transdermal nicotine absorption from nicotine-polluted surfaces might lead to an overestimation of cotinine-based ETS exposure.

Biomarkers for potentially genotoxic compounds related to ETS exposure

In contrast to biomonitoring for surrogate markers, which can give only indirect evidence of the uptake of substances of toxicological relevance, biomonitoring for potentially genotoxic compounds directly reflects the internal dose of substances possibly implicated in the process of carcinogenesis. However, with the exception of tobacco-specific nitrosamines, the genotoxic substances found in tobacco smoke occur almost ubiquitously in the environment and are taken up from various sources. Hence, for evaluating the contribution of ETS exposure to the total body burden, it is important to assess the background levels of these biomarkers. ETS exposure-related biomonitoring data for specific genotoxic compounds such as benzene, polycyclic

aromatic hydrocarbons (PAH), aromatic amines and tobacco-specific nitrosamines (TSNA) as well as group-selective analyses for various DNA adducts, urinary thioethers and mutagenic activity are available (Table 2).

Benzene

Benzene occurs ubiquitously in the environment, with traffic exhaust being the most important source.¹⁹ Average benzene concentrations in rural and urban environments of 1–10 and 10–20 $\mu\text{g}/\text{m}^3$, respectively, have been reported in Germany.^{20,21} Cigarettes were found to emit 30–50 $\mu\text{g}/\text{cigarette}$ in MS and 345–653 $\mu\text{g}/\text{cigarette}$ of benzene in SS.²² In a household survey including 230 homes in Germany, the median benzene concentrations were 6.9 $\mu\text{g}/\text{m}^3$ in households with nonsmokers and 9.3 $\mu\text{g}/\text{m}^3$ in homes with at least one smoker.²³ Corresponding levels reported for the USA were 7 and 10.5 $\mu\text{g}/\text{m}^3$, respectively.²⁴ Various biomarkers can be used for the biological monitoring of benzene exposure.²⁵ Determination of benzene in blood²⁶ and exhalate²⁷ is suitable for biomonitoring acute exposure. Due to the longer half-lives of 6–12 h, the urinary benzene metabolites *trans,trans*-muconic acid and phenylmercapturic acid should preferably be used in field studies.^{28,29} However,

trans,trans-muconic acid is also a metabolite of the food preservative sorbic acid, and thus may not be a specific biomarker for low-level environmental benzene exposure.³⁰ Investigations under real-life conditions showed no, or only a marginal contribution, of ETS exposure to the background of benzene in nonsmokers (Table 2). Analysis of variance in one study showed that at most 15 % of the variation is explained by ETS exposure.³¹ This is in good agreement with an air monitoring survey in 49 homes, which apportioned 11% of indoor air benzene to ETS.³² Taken together, the biomonitoring data show that ETS exposure is only a minor source of the total benzene burden.

Polycyclic aromatic hydrocarbons (PAH)

PAH are formed during incomplete combustion of organic materials and are ubiquitously distributed in the environment. The main source of intake of PAH is the diet. In particular, fried, grilled, smoked and cured foods as well as leafy vegetables contain PAH mostly in the upper p.p.b. range with benzo[a]pyrene occurring in the lower p.p.b. range.³³ It is estimated that >90 % of the total PAH body burden originates from the diet.^{32,34} In the MS and SS of a cigarettes about 10 and 100 ng of benzo[a]pyrene, respectively, are emitted.³⁴ Indoor air concentrations of benzo[a]pyrene were found to

Table 2 Biomonitoring for exposure to genotoxic compounds*

Precursor in ETS	Other sources	Biomarker	Significant increase after real-life ETS exposure
Benzene	Traffic exhaust, combustion, fuels	Benzene in blood or exhalate	No ^{11a} /Yes ^{9,11b}
PAH	Sorbic acid Combustion, diet, ambient air	1,1-MA in urine DNA adducts in WBC Albumin adducts 1-Hydroxypyrene in urine	(Yes ¹⁷) ^b No ¹⁹ /Yes ¹⁸ No ³ /Scherer, in preparation
4-ABP	Gas or kerosene burners, diesel exhaust	4-ABP-haemoglobin adducts	No ^{4b,29} /Yes ^{44,31c}
NNK	—	NNAL and NNAL-glucuronide in urine	(Yes ⁵⁵) ^c
NNK, NNN	—	HBP-haemoglobin adducts	No ^{4b}
Polycyclic aromatics	Diet, ambient air, endogenous	bulky DNA adducts in WBC and placenta	No ^{64,65,62,68} /(Yes ⁶⁶) ^c
NDMA, NNK, (NDEA)	Diet, ambient air, endogenous formation	3-Methyl-/3-Ethyl-adenine in urine	(No ⁷²) ^c
?	Oxidative stress (exogenous and endogenous)	8-OHdG in placenta	No ⁶⁸
Carbonyls (acrolein)	Diet	Thioethers in urine	No ⁷³
Aromatic amines(?)	Diet	Mutagenic activity in urine	No ⁷⁴ – ⁷⁶ /Yes ⁷⁹

*Abbreviations: 1,1-MA: *trans,trans*-muconic acid; PAH: polycyclic aromatic hydrocarbons; WBC: white blood cells; 4-ABP: 4-aminobiphenyl; NNK: 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNAL: 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; NNN: N-nitrosodimethylaniline; HBP: 4-hydroxy-1-(3-pyridyl)-1-butanone; NDMA: N-nitrosodimethylaniline; NDEA: N-nitrosodiethylaniline; 8-OHdG: 8-hydroxy-2'-deoxyguanosine. ^bSignificant correlation between score for ETS exposure and DNA adducts. (Only 4 of 21 samples had detectable adduct levels.) ^cAfter high experimental exposure to sidestream smoke for 3 h. ^dNo systematic assessment of ETS exposure.

^eAfter high experimental exposure to ETS for 8 h.

be 5 ng/m³ in homes with smokers and 3 ng/m³ in homes with nonsmokers.¹⁵ The ultimate carcinogenic metabolites of benzo[a]pyrene and other PAH can bind to DNA or proteins. The long life-time of erythrocytes (120 days) leads to the accumulation of haemoglobin adducts and makes them suitable biomarkers reflecting exposure during the last 4 months. Because of the shorter half-life of albumin (20–24 days), its adducts reflect more recent exposure. Protein adducts can be taken as an index of the biologically effective dose of a carcinogen. Urinary excretion of monohydroxy phenanthrenes and 1-hydroxypyrene are also used for biomonitoring PAH exposure by passive smokers.¹⁶ In most investigations, no increase in PAH biomarkers was found after ETS exposure. The two studies^{17,18} showing significant ETS-related increases (Table 2) remain doubtful in view of the fact that (a) other working groups found only a small, if any, difference in protein PAH adduct levels between smokers and nonsmokers^{19–21}, (b) even extremely high exposure to ETS did not increase the urinary excretion of PAH metabolites²² and (c) the estimated intake of PAH suggested an overwhelming role of diet.²³ Summarizing the available evidence for PAH exposure by passive smoking, it can be stated that the high background exposure to PAH from other sources, particularly from diet, excludes biomonitoring of these substances from being of any value in ETS risk assessment.

Aromatic amines

Haemoglobin adducts of 4-aminobiphenyl (4-ABP) and 3-aminobiphenyl (3-ABP) have been used to determine exposure to these substances by tobacco smoke. MS yields of 4-ABP and 3-ABP were found to be 2.4–4.6 and 2.7–5.0 ng/cigarette, respectively.^{42,43} SS was found to emit 143 and 132 ng/cigarette of 4-ABP and 3-ABP, respectively.⁶ In an experimental room containing an ETS-related carbon monoxide concentration of 8 p.p.m., a level of about 5 ng/m³ of 4-ABP was measured (Grimmer, personal communication). As yet, no other sources for 4- and 3-ABP have been identified, apart from the fact that 4-ABP was used in the dye industry decades ago.⁴⁴ However, 3- and 4-nitrobiphenyl, which are emitted by kerosene heaters and gas burners,⁴⁵ as well as by diesel engines, form the same haemoglobin adduct as do 3- and 4-ABP and must, therefore, be considered confounding factors. Controversial results on an effect of ETS exposure on the 4-ABP haemoglobin adduct level were reported (Table 2). In the most extensive study,⁴⁶ no ETS-related increase was observed. Nonsmokers living in cities were found to have 4-ABP-haemoglobin adduct levels 10–30 pg/g higher than nonsmokers living in rural areas.^{47,48} This suggests that sources other than ETS are also important.

Tobacco-specific nitrosamines (TSNA)

TSNA are mainly formed from the tobacco alkaloids nicotine, nornicotine, anabasine and anatabine during fermentation of tobacco leaves. 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and N-nitrosornicotine (NNN) are the biologically most important TSNA.⁴⁹ MS yields of commercial cigarettes were reported to be 17–306 ng/cigarette for NNK and 34–675 ng/cigarette for NNN.⁵⁰ Corresponding SS yields were 180–671 ng/cigarette and 141–348 ng/cigarette, respectively.⁵⁰ Reported concentrations of TSNA in rooms where smoking occurs, range from 0.2 to 29.3 ng/m³ for NNK and 0.7–23 ng/m³ for NNN.^{51–53} As biomarkers for NNK exposure, urinary NNAL, the corresponding alcohol of NNK, and its glucuronide as well as pyridyloxobutylated haemoglobin and DNA have been used.⁵⁴ DNA and haemoglobin adducts, which also indicate exposure to NNN, release 4-hydroxy-1-(3-pyridyl)-1-butanone (HPB) upon hydrolysis. There is no systematic investigation on the influence of everyday ETS exposure on urinary excretion of NNAL and NNAL glucuronide. In an experimental study, Hecht *et al.*⁵⁵ exposed five nonsmokers for 3 h in a small chamber of 16 m³ air volume to the extremely high SS concentrations of up to 230 µg/m³ nicotine and up to 263 ng/m³ NNK. Under these conditions, the urinary excretion of free and conjugated NNAL increased from 31 ± 41 pmol/d before exposure to 127 ± 74 pmol/d after exposure. The authors state that NNK uptake in smokers is about 120 times higher than in nonsmokers exposed to sidestream smoke.⁵⁶ We found in four of nine nonsmokers detectable NNAL and NNAL glucuronide levels (detection limit about 5 pmol/l) with an average excretion rate of 41 ± 52 pmol/d.⁵⁶ These results are comparable to the findings with nonsmokers before experimental SS exposure in the study of Hecht *et al.*⁵⁵ and suggest that the ETS-related NNK exposure under real-life conditions amounts to about 1% of the NNK dose in smokers.

HPB-releasing haemoglobin adducts could be tobacco-specific biomarkers of chronic exposure to NNK and NNN. Mean adduct levels for smokers and nonsmokers were 80 and 29 fmol/g haemoglobin, respectively, as reported by the working group of Hecht⁵⁷ and 69 and 34 fmol/g haemoglobin as found by the working group of Richter.⁴⁷ There was a large overlap between smokers and nonsmokers in both studies. The small differences in the HPB-adducts found between smokers and nonsmokers are remarkable. According to the NNK and NNN levels in MS and ETS as well as the urinary NNAL excretion rates discussed above, smokers should have HPB adduct levels at least two orders of magnitude higher than ETS-exposed nonsmokers. The lack of such a large difference might be explained by induction of TSNA detoxification and/or inhibition of TSNA activation in smokers.

Evidence for the latter hypothesis is provided by a recent finding suggesting that nicotine in smoking doses can inhibit NNK activation in rats.⁶⁴ In pregnant women, however, no relationship between ETS exposure (either based on self-reported exposure or classified by urinary cotinine levels) and HPB-haemoglobin adduct levels was observed.⁶⁵ In theory, NNK may also be formed endogenously by nitrosation of nicotine. No evidence for nitrosation of nicotine or cotinine to the possible nitrosation product 4-(*N*-methylnitrosamino)-4-(3-pyridyl)butyric acid (iso-NNAC) was observed.⁶⁶ In addition, the nitrosoproline test, which reveals an increased endogenous nitrosation capacity of smokers,⁶⁷ did not show elevated nitrosation in nonsmokers experimentally exposed to high ETS doses.^{68,69} The two biomarkers for TSNA exposure, namely urinary NNAL and the HPB-haemoglobin adducts, need further validation before conclusions on ETS-related TSNA doses and possibly implicated risks can be drawn.

Various DNA adducts

Cigarette smoke condensate (CSC), when dermally applied to mice, was found to form various, as yet unidentified DNA adducts detectable by the "P-postlabelling method."⁷⁰ In human biomonitoring studies, available tissues and cells are limited to leucocytes, oral mucosa, exfoliated epithelial bladder cells, bronchial lavage cells and placenta. We are aware of only a few investigations where experimental^{64,65} or real-life exposure to ETS⁶⁶⁻⁶⁹ has been considered causative for the formation of DNA adducts detectable by the "P-postlabelling method. In one study, it was suggested that passive smoking might lead to the formation of a tobacco smoke-related adduct with placental DNA.⁶⁶ However, the evidence was weak, since ETS exposure was only reported for those nonsmokers who had detectable levels of this adduct. In another study with systematic assessment of ETS exposure in pregnant women, no increase in placental DNA adducts in passive or active smokers was found.⁶⁷ No additional DNA adducts or increases in DNA adduct levels after ETS exposure were observed in peripheral monocytes,⁶⁴ lymphocytes⁶⁵ or white blood cells.⁶⁷

In addition to NNK, *N*-nitrosodimethylamine (NDMA) is another methylating agent in tobacco smoke. MS and SS yields were reported to be 0.1–20 and 143–1040 ng/cigarette, respectively.⁷¹ ETS concentrations of NDMA in rooms with moderate smoking were 20–50 ng/m³.⁶⁹ *N*-Nitrosodiethylamine (NDEA), *N*-nitrosomethylethylamine (NMEA) and ethylhalogenides are potential ethylating agents in tobacco smoke. However, only trace amounts of these nitrosamines are detectable in tobacco smoke,⁷⁰ whereas about 1 µg of ethylchloride was found in MS.⁷¹ In an experimental study, no

increase in the urinary excretion of 3-methyladenine or 3-ethyladenine was observed in nonsmokers exposed to high ETS concentrations.⁷²

The promutagenic DNA adduct 8-hydroxy-2'-deoxyguanosine (8-OHdG) formed by reactive oxygen species is an established biomarker for oxidative DNA damage.⁷³ *In vitro* experiments have shown that the MS of cigarettes can induce 8-OHdG adducts after metabolic activation, the gaseous phase being responsible for this effect.⁷⁴ In a study aimed at systematically assessing ETS exposure in pregnant women, no ETS exposure-related increases in placental 8-OHdG adduct levels were found.⁶⁸

Electrophilic compounds

Electrophilic chemicals are considered potential toxicants, mutagens and/or carcinogens because they may covalently bind to cellular macromolecules such as proteins, RNA and DNA.⁷⁵ Conjugation of electrophiles to glutathione (GSH), either spontaneously or enzymatically by means of glutathione *S*-transferases (GST), in most cases indicates a detoxification process. The GSH adducts are further metabolized to form *S*-substituted *N*-acetyl-L-cysteine conjugates (thioethers, mercapturic acids), which are renally excreted. Urinary thioethers have been used as group-selective biomarkers for the exposure to electrophiles.⁷⁶ Passive smoking under real-life conditions did not lead to a measurable increase in urinary thioether excretion (Table 2).⁷⁶ In an experimental study under dietary control and exposure to high ETS concentrations over a period of 8 h, a significant increase in excretion as compared to sham exposure was observed.⁶⁵ This effect was found to be caused by exposure to the gaseous phase of ETS. As a specific thioether, 3-hydroxypropyl mercapturic acid was identified, which is probably related to acrolein exposure.^{65,74}

Mutagens

The mutagenic properties of CSC in short-term tests after metabolic activation are well established.⁷⁷ Measurements of the urinary mutagenicity in nonsmokers experimentally exposed to high doses of ETS yielded controversial results.⁷⁸ The influence of real-life ETS exposure on urinary mutagenicity has been determined in five populations.⁷⁶⁻⁷⁹ No significant influence of ETS exposure was found in all but one investigation.⁷⁹ More extensive studies are needed in order to quantify the contribution of ETS exposure to urinary mutagenicity, which is primarily dominated by dietary factors.⁸⁰

Markers for biological effects related to ETS exposure

By definition, biological effect markers are not specific for the underlying exposure but reflect

physiological or toxicological responses to a great number of factors. Therefore, it is extremely important to control for confounding factors when applying these markers in population studies. Data about various biological effect markers in relation to ETS exposure are summarized in Table 3.

Cytogenetic effects

Tobacco smoke and its condensate have been shown to induce various cytogenetic damages and effects such as chromosomal aberrations (CA), sister chromatid exchanges (SCE) and micronuclei (MN) *in vitro* and *in vivo*.⁷⁷⁻⁸¹ The influence of ETS exposure on cytogenetic parameters has been investigated in three limited studies^{77,82,83} and one more extensive approach.⁸⁴ No effect of passive smoking was found, probably because these tests were not sensitive enough to detect effects arising from low exposures, such as passive smoking.

Aryl hydrocarbon hydroxylase (AHH) induction

It is well documented that cigarette smoking can induce, via the Ah receptor, enzymes involved in the activation (e.g. cytochrome P4501A1) and detoxification (glutathione S-transferase, UDP-glucuronyltransferase) of xenobiotics. Induction is particularly high in the placenta.⁸⁵ The role of exposure to ETS in AHH induction has been investigated in human placenta at the enzyme^{86,87} and messenger RNA level.⁸⁸ The results suggest that passive smoking may exert an inducing effect in the placenta. Further investigations are needed to clarify this possible effect of ETS exposure. Whether induction of the AHH enzyme system may increase the risk after exposure to PAH or other chemicals which are activated by cytochrome P450 enzymes or whether induction may be protective, as suggested by Remmer,⁸⁹ is an open question.

Urinary hydroxyproline (HOP) excretion

Increased urinary excretion of HOP is an established biomarker for certain osteopathic destructions, some endocrinological disorders and severe burns.⁹⁰ Since HOP is a degradation product of lung collagen and elastin induced by exposure to nitrogen dioxide (NO₂), this marker has also been used to study the effect of low NO₂ exposures arising from environmental automobile exhausts as well as active and passive smoking.⁹¹ The results reported in the literature on the effects of active and passive smoking on urinary HOP excretion are controversial: while Kasuga⁹² found significantly increased levels of this marker in smokers and passive smokers, these findings were not confirmed by others.⁹³⁻⁹⁵ In a recent study, no relationship between personal NO₂ and ETS exposure and urinary excretion of HOP or desmosine, a catabolic product of elastin, was found.⁹⁶

Effect markers for cardiovascular diseases

The pathomechanism by which active or passive smoking might increase the risk of cardiovascular disease is not completely understood. Possible effects include reduced oxygen supply, endothelial injury, reduction in high-density lipoprotein (HDL) cholesterol, increase in low-density lipoprotein (LDL) cholesterol, increased LDL oxidation with subsequent foam cell formation, increased platelet activation and coagulation.⁹⁷⁻⁹⁹ Physiological changes in parameters related to these effects can be used as biological effect markers. In a couple of studies, the influence of ETS exposure under experimental and real-life conditions on possible effect markers has been investigated (Table 3).

In a small ETS field study, no effect of ETS exposure on either urinary 2,3-dinor-thromboxane B₂ or 2,6-dinor-6-ketoprostaglandin F_{1α} excretion was found.¹⁰⁰ Sinzinger and coworkers reported a significant reduction in platelet sensitivity to

Table 3 Monitoring of biological effects related to ETS exposure^a

Biological effect or biomarker	ETS component possibly implicated	Significant effect of ETS exposure under real-life conditions
SCE, CA in lymphocytes	ETS particulate matter (?)	No ^{77,82-84}
AHH induction in placenta	PAH, others	Yes ⁸⁶⁻⁸⁸
Hydroxyproline in urine	NO ₂ (?)	No ^{92,93} /Yes ⁹¹
Total cholesterol	?	No ^{109,110} /Yes ¹¹⁹
HDL in plasma	?	No ^{109,110} /Yes ^{107,119}
LDL in plasma	?	No ^{109,110,119}
Platelet aggregation	?	Yes ¹⁰²
Fibrinogen in plasma	?	Yes ¹⁰⁵
Tx-M and PGI-M in urine	?	No ⁹⁸
Carotid wall thickness	?	Yes ^{103,104}

^aAbbreviations: SCE: sister chromatid exchanges; CA: chromosomal aberrations; AHH: aryl hydrocarbon hydroxylase; PAH: polycyclic aromatic hydrocarbons; HDL: high-density lipoprotein; LDL: low-density lipoprotein; Tx-M: 2,3-dinor-thromboxane B₂; PGI-M: 2,6-dinor-6-ketoprostaglandin F_{1α}.

In two large population studies, ETS exposure was found to significantly increase the carotid wall thickness^{101, 101} and the plasma fibrinogen concentration.¹⁰⁵ Although the observed changes are small and the biological significance in terms of cardiovascular risk is unclear, the effects are relatively large in relation to the effect observed in active smokers. Both studies claim to have controlled for the major cardiovascular risk factors such as age, body mass index, ethanol intake, hypertension, diabetes, and total fat intake. However, since it is well established that the passive smoking status can be associated with an unfavourable constellation of a great number of cardiovascular risk factors,¹⁰⁶ it seems doubtful whether statistical control of confounding factors can be complete.

During the past 10–20 years, a large amount of data on biomonitoring of ETS exposure has accumulated. Cotinine in body fluids of nonsmokers has proven to be an acceptable, although nonideal biomarker for assessing ETS exposure dating back 1–3 days. This marker has been and will be almost routinely used in field studies in addition to or for validation of self-reported exposure to ETS. Nonsmokers, chronically exposed to ETS under real-life conditions, show cotinine concentrations in blood, saliva and urine at least two orders of magnitude lower than those of current cigarette smokers. Although nicotine uptake by active and passive smoking differs in many aspects, this comparison gives a good estimate for the dose difference between smoking and ETS exposure. COHb and CO in the exhalate as well as thiocyanate in body fluids, which all are also used as biomarkers for ETS exposure, are less suitable due to their low

Biomonitoring for exposure to ETS-related genotoxic compounds might provide more relevant information for risk assessment than do the surrogate biomarkers cotinine, COHb and thiocyanate. Data are available for exposure to benzene, PAH, aromatic amines and tobacco-specific nitrosamines. In addition, various DNA adducts in nucleated blood cells and placenta as well as urinary thioether excretion and mutagenic activity have been investigated in relation to ETS exposure. The results obtained are not conclusive, but suggest that every-day ETS exposure, if at all, only marginally increases the levels of these markers above background levels. Three major difficulties have to be considered when interpreting these data: (1) it is yet unknown which carcinogens in tobacco smoke are responsible for tumor induction in humans. As a consequence, it is not possible to focus the biological monitoring on the appropriate chemicals or classes of chemicals. (2) Population biomonitoring is limited to readily available body fluids (blood, urine, saliva), cells (blood cells, oral mucosa cells, exfoliated bladder cells) and tissues (skin, nails, hair, placenta). With respect to the most widely discussed cancer risk attributed to ETS exposure, namely lung cancer, these materials represent no target cells or tissues, but allow only the determination of surrogate biomarkers. (3) With the exception of TSNA, the biomarkers for exposure to genotoxic compounds applied today are not related to substances unique for tobacco smoke, but occur ubiquitously in the environment and in food. Therefore, the existing background exposure to these substances has to be considered when evaluating an ETS-related increase in risk. As discussed above, the most promising biomarkers for TSNA exposure, i.e. urinary NNAL and the TSNA-related haemoglobin adducts, need further validation before being applicable for the biomonitoring of ETS exposure.

ETS exposure was not found to exert cytogenetic effects such as SCE or CA in peripheral lymphocytes. However, these biomarkers may not be sensitive enough to indicate effects after low exposure, such as passive smoking.

Controversial results were reported on the effect of ETS exposure on urinary hydroxyproline (HOP) excretion. The findings of Kasuga,²³ who found a significant dose-related effect of ETS exposure on urinary HCP excretion, could not be confirmed by other investigators.²²⁻²⁴ Further studies are needed to clear this discrepancy.

A small AHH-inducing effect of ETS exposure in human placenta was found. The biological significance of this finding is unclear, because induction

comprises most probably enzymes involved in both activating and detoxifying pathways of carcinogens. It would be of interest to investigate ETS exposure-related enzyme induction in other relevant organ systems (e.g. the lung).

Significant, unfavourable effects of ETS exposure were reported for some biomarkers, which are regarded to be indicative for the development of cardiovascular diseases. These include total serum cholesterol, HDL, plasma fibrinogen, platelet aggregation and carotid wall thickness. The extent of

these effects was surprisingly high as compared to that observed with smokers. To explain this discrepancy, it is hypothesized that nonsmokers are more sensitive to some tobacco smoke components than are smokers.^{101,102} An alternative explanation would be that passive smokers differ from nonsmokers not only in their exposure to ETS, but also in a series of other cardiovascular risk factors.¹⁰⁰ The alleged detrimental effects of ETS exposure on the cardiovascular system are an important issue for future research.

References

- 1 Environmental Protection Agency. *Respiratory Health Effects of Passive Smoking: Lung Cancer and other disorders*. Washington, EPA 600/6-90/006F, 1992.
- 2 Jarvis MJ. Application of biochemical intake markers to passive smoking measurement and risk estimation. *Mutat Res* 1989; **222**: 101–110.
- 3 Coburn RF. Endogenous carbon monoxide production. *New Engl J Med* 1970; **282**: 207–209.
- 4 Guerin MR, Jenkins RA, Tomkins BA. *The Chemistry of Environmental Tobacco Smoke: Composition and Measurement*. Boca Raton, Ann Arbor, London, Tokyo, Lewis Publishers, 1992.
- 5 Borgers D, Junge B. Thiocyanate as an indicator of tobacco smoking. *Prev Med* 1979; **8**: 351–357.
- 6 Klus H, Kuhn H. Verteilung verschiedener Tabakrauchbestandteile auf Haupt- und Nebstromrauch (Eine Übersicht). *Beitr Tabakforsch Int* 1982; **11**: 229–265.
- 7 Baker RR, Proctor CJ. The origins and properties of environmental tobacco smoke. *Environ Int* 1990; **16**: 231–245.
- 8 Phillips K, Howard DA, Browne D, Lowsley JM. Assessment of personal exposures to environmental tobacco smoke in British nonsmokers. *Environ Int* 1994; **20**: 693–712.
- 9 Scherer G *et al.* Contribution of tobacco smoke to environmental benzene exposure in Germany. *Environ Int* 1995; **21**: 779–789.
- 10 Domino E-F, Hornbach E, Demana T. Relevance of nicotine content of common vegetables to the identification of passive tobacco smokers. *Med Sci Res* 1993; **21**: 571–572.
- 11 Scherer G *et al.* Pharmacokinetics of nicotine, cotinine, and 3'-hydroxycotinine in cigarette smokers. *Klin Wochenschr* 1988; **66**, Suppl: 5–11.
- 12 Pirkle JL *et al.* Exposure of the US population to environmental tobacco smoke. *JAMA* 1996; **275**: 1233–1240.
- 13 Reasor MJ, Will JA. Assessing exposure to environmental tobacco smoke: is it valid to extrapolate from active smoking? *J Smoking-Related Dis* 1991; **2**: 111–127.
- 14 Scherer G *et al.* Importance of exposure to gaseous and particulate phase components of tobacco smoke in active and passive smokers. *Int Arch Occup Environ Health* 1990; **62**: 459–466.
- 15 Haley NJ, Sepkovic DW, Hoffmann D. Elimination of cotinine from body fluids: Disposition in smokers and nonsmokers. *Am J Publ Health* 1989; **79**: 1046–1048.
- 16 Benowitz NL. Cotinine as a biomarker of environmental tobacco smoke exposure. *Epidemiol Rev* 1996; **18**: 188–204.
- 17 Neurath GB *et al.* Gas-particulate phase distribution and decay rates of constituents in ageing environmental tobacco smoke. *Environ Technol* 1991; **12**: 581–590.
- 18 Ogden MW. Occupational exposure to environmental tobacco smoke. *JAMA* 1996; **14**: 441.
- 19 Fishbein L. Benzene. Uses, occurrences and exposure. In Fishbein L, O'Neill IK (eds): *IARC Scientific Publications No. 85: Environmental Carcinogens Methods of Analysis and Exposure Measurement, Volume 10 - Benzene and Alkylated Benzenes*. Lyon, International Agency for Research on Cancer, 1988: pp.67–96.
- 20 Müller J. Innen- und Außenluftmessungen an einer innerstädtischen Hauptverkehrsstraße. *Staub-Reinhalt Luft* 1991; **51**: 147–154.
- 21 Chriske HW, Brockhaus A, Ewers U. Benzolbelastung von im Außen- und Innendienst tätigen Personen einer verkehrsreichen Großstadt. *Arbeitsmed Sozialmed Präventivmed* 1991; **26**: 483–485.
- 22 Brunnemann KD, Kagan MR, Cox JE, Hoffmann D. Analysis of 1,3-butadiene and other selected gas-phase components in cigarette mainstream and sidestream smoke by gas chromatography-mass selective detection. *Carcinogenesis* 1990; **11**: 1863–1868.
- 23 Krause C *et al.* Occurrence of volatile organic compounds in the air of 500 homes in the Federal Republic of Germany. In Seifert B, Esdorn H, Fischer M, Rüden H, Wegner J (eds): *Indoor Air '87, Proceedings of the 4th International Conference on Indoor Air Quality and Climate, Volatile Organic Compounds, Combustion Gases, Particles and Fibres, Microbiological Agents*. Berlin, Institute for Water, Soil and Air Hygiene, 1987: vol 1, pp.102–111.
- 24 Wallace LA. Major sources of benzene exposure. *Environ Health Perspect* 1989; **82**: 165–169.
- 25 Lauwerys R, Buchet JP. Biological monitoring of exposure to benzene, toluene and xylene. In Fishbein L, O'Neill IK (eds): *Environmental Carcinogens Methods of Analysis and Exposure Measurement, Volume 10 - Benzene and Alkylated Benzenes*. Lyon, International Agency for Research on Cancer, 1988: pp.205–222.

- 26 Angerer J, Scherer G, Schaller KH, Müller J. The determination of benzene in human blood as an indicator of environmental exposure to volatile aromatic compounds. *Fresenius J Anal Chem* 1991; **339**: 740-742.
- 27 Riedel K *et al.* Determination of benzene and alkylated benzenes in ambient and exhaled air by microwave desorption coupled with gas chromatography-mass spectrometry. *J Chromatogr* 1996; **719**: 383-389.
- 28 Inoue O *et al.* Urinary t,t-muconic acid as an indicator of exposure to benzene. *Br J Ind Med* 1989; **46**: 122-127.
- 29 Strommel P *et al.* Determination of S-phenylmercapturic acid in the urine - an improvement in the biological monitoring of benzene exposure. *Carcinogenesis* 1989; **10**: 279-282.
- 30 Ducos P *et al.* trans, trans-Muconic acid, a reliable biological indicator for the detection of individual benzene exposure down to the ppm level. *Int Arch Occup Environ Health* 1992; **64**: 309-313.
- 31 Heavner DL, Morgan WT, Ogden MW. Determination of volatile organic compounds and ETS apportionment in 49 homes. *Environ Int* 1995; **21**: 3-21.
- 32 Hattmer-Frey HA, Travis CC. Benzo-a-pyrene: Environmental partitioning and human exposure. *Toxicol Ind Health* 1991; **7**: 141-157.
- 33 World Health Organization. *Guidelines for drinking-water quality. Vol. 1 Recommendations*. Geneva, World Health Organization, 1984.
- 34 Grimmer G, Naujack KW, Dettbarn G. Gaschromatographic determination of polycyclic aromatic hydrocarbons, aza-arenes, aromatic amines in the particle and vapor phase of mainstream and sidestream smoke of cigarettes. *Toxicol Lett* 1987; **35**: 117-124.
- 35 Chuang JC, Mack GA, Kuhlman MR, Wilson NK. Polycyclic aromatic hydrocarbons and their derivatives in indoor and outdoor air in an eight-home study. *Atmospheric Environment* 1991; **25**: 369-380.
- 36 Martin F *et al.* Urinary excretion of hydroxy-phenanthrenes after intake of polycyclic aromatic hydrocarbons. *Environ Int* 1989; **15**: 41-47.
- 37 Perera FP *et al.* DNA adducts, protein adducts, and sister chromatid exchange in cigarette smokers and nonsmokers. *J Natl Cancer Inst* 1987; **79**: 449-456.
- 38 Crawford FG *et al.* Biomarkers of environmental tobacco smoke in preschool children and their mothers. *J Natl Cancer Inst* 1994; **86**: 1398-1402.
- 39 Autrup H, Beck Vestergaard A, Okkels H. Transplacental transfer of environmental genotoxins: polycyclic aromatic hydrocarbon-albumin in non-smoking women, and the effect of maternal GSTM1 genotype. *Carcinogenesis* 1995; **16**: 1305-1309.
- 40 Pastorelli R *et al.* Hemoglobin adducts of benzo[a]pyrene diol epoxide in newspaper vendors: association with traffic exhaust. *Carcinogenesis* 1996; **17**: 2389-2394.
- 41 Tas S, Buchet JP, Lauwerys R. Determinants of benzo[a]pyrene diol epoxide adducts to albumin in workers exposed to polycyclic aromatic hydrocarbons. *Int Arch Occup Environ Health* 1994; **66**: 343-348.
- 42 International Agency for Research on Cancer. *Tobacco Smoking. IARC Monographs on the evaluation of the carcinogenic risk of chemicals to humans*. Lyon, IARC, 1986: vol 38.
- 43 Grimmer G, Schneider D. Intercept-reactant method for the determination of aromatic amines in mainstream tobacco smoke. *Beitr Tabakforsch Int* 1995; **16**: 141-156.
- 44 Hammond SK *et al.* Relationship between environmental tobacco smoke exposure and carcinogen-hemoglobin adduct levels in nonsmokers. *J Natl Cancer Inst* 1993; **85**: 474-478.
- 45 Tokiwa H, Nakagawa R, Horikawa K. Mutagenic/carcinogenic agents in indoor air pollutants: The dinitropyrenes generated by kerosene heaters and fuel gas and liquefied petroleum gas burners. *Mutat Res* 1985; **157**: 39-47.
- 46 Richter E *et al.* Comparison of biomarkers for exposure to environmental tobacco smoke: Cotinine and haemoglobin adducts from aromatic amines and tobacco-specific nitrosamines in pregnant smoking and nonsmoking women, in Maroni M (ed): *healthy buildings '95 an international conference on healthy buildings in mild climate*. Milano, healthy buildings '95, 1995: pp. 599-604.
- 47 Falter B, Kutzer C, Richter E. Biomonitoring of hemoglobin adducts: aromatic amines and tobacco-specific nitrosamines. *Clin Invest* 1994; **72**: 364-371.
- 48 Richter E. Gesundheitsgefährdung durch Passivrauchen. *Deutsche Gesellschaft für experimentelle und klinische Pharmakologie und Toxikologie (DGPT) Mitteilungen* 1996; **19**: 65-66.
- 49 Hoffmann D, Rivenson A, Hecht SS. The biological significance of tobacco-specific N-nitrosamines: Smoking and adenocarcinoma of the lung. *Crit Rev Toxicol* 1996; **26**: 199-211.
- 50 Hoffmann D, Brunnemann KD, Prokopczyk B, Djordjevic MV. Tobacco-specific N-nitrosamines and Areca-derived N-nitrosamines: Chemistry, biochemistry, carcinogenicity, and relevance to humans. *J Toxicol Environ Health* 1994; **41**: 1-52.
- 51 Brunnemann KD, Cox JE, Hoffmann D. Analysis of tobacco-specific N-nitrosamines in indoor air. *Carcinogenesis* 1992; **13**: 2415-2418.
- 52 Klus H *et al.* Tobacco-specific and volatile N-nitrosamines in environmental tobacco smoke of offices. *Indoor Environment* 1992; **1**: 348-350.
- 53 Tricker AR *et al.* Tobacco-specific and volatile nitrosamines in indoor air of smoker and nonsmoker homes, in Weber L (ed): *Indoor Air Pollution. Proceedings of the International Conference held at the University of Ulm, Ulm, Germany, 5-7 October 1994*. Rothenfluh, Switzerland, Indoor Air International, 1994: pp. 310-316.
- 54 Hecht SS. Recent studies on mechanisms of bioactivation and detoxification of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), a tobacco-specific lung carcinogen. *Crit Rev Toxicol* 1996; **26**: 163-181.
- 55 Hecht SS *et al.* A tobacco-specific lung carcinogen in the urine of men exposed to cigarette smoke. *New Engl J Med* 1993; **329**: 1543-1546.

- 56 Meger M *et al.* Metabolites of 4-(N-methylnitrosamino)-1-(3-pyridyl)-1-butanone in urine of smokers. *European Journal of Cancer Prevention* 1996; 5 (Suppl. 1): 121-124.
- 57 Hecht SS *et al.* Tobacco-specific nitrosamine adducts: Studies in laboratory animals and humans. *Environ Health Perspect* 1993; 99: 57-63.
- 58 Richter E, Tricker AR. Nicotine inhibits the metabolic activation of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in rats. *Carcinogenesis* 1994; 15: 1061-1064.
- 59 Tricker AR *et al.* Evaluation of 4-(N-methylnitrosamino)-4-(3-pyridyl)butyric acid as a potential monitor of endogenous nitrosation of nicotine and its metabolites. *Carcinogenesis* 1993; 14: 1409-1414.
- 60 Hoffmann D, Brunnemann K. Endogenous formation of N-nitrosoproline in cigarette smokers. *Cancer Res* 1983; 43: 5570-5574.
- 61 Hoffmann D, Haley NJ, Adams JD, Brunnemann KD. Tobacco sidestream smoke: Uptake by nonsmokers. *Prev Med* 1984; 13: 608-617.
- 62 Scherer G, Adlkofer F. Endogenous formation of N-nitrosoproline in smokers and nonsmokers. *Banbury Report* 1986; 23: 137-148.
- 63 Randerath E, Mittal D, Randerath K. Tissue distribution of covalent DNA damage in mice treated dermally with cigarette 'tar': preference for lung and heart DNA. *Carcinogenesis* 1988; 9: 75-80.
- 64 Holz O *et al.* ³²P-postlabelling analysis of DNA adducts in monocytes of smokers and passive smokers. *Int Arch Occup Environ Health* 1990; 62: 299-303.
- 65 Scherer G, Conze C, Tricker AR, Adlkofer F. Uptake of tobacco smoke constituents on exposure to environmental tobacco smoke (ETS). *Clin Invest* 1992; 70: 352-367.
- 66 Everson RB *et al.* Detection of smoking-related covalent DNA adducts in human placenta. *Science* 1986; 231: 54-57.
- 67 Binkova B *et al.* DNA adducts and personal air monitoring of carcinogenic polycyclic aromatic hydrocarbons in an environmentally exposed population. *Carcinogenesis* 1995; 16: 1037-1046.
- 68 Daube H *et al.* DNA adducts in human placenta in relation to tobacco smoke exposure and plasma antioxidant status. *J Cancer Res Clin Oncol* 1997; 123: 141-151.
- 69 Stehlik G, Richter O, Altmann H. Concentration of dimethylnitrosamine in the air of smoke-filled rooms. *Ecotoxicol Environ Safety* 1982; 6: 495-500.
- 70 Tricker AR, Ditrach C, Preussmann R. N-Nitroso compounds in cigarette tobacco and their occurrence in mainstream tobacco smoke. *Carcinogenesis* 1991; 12: 257-261.
- 71 Jenkins RW *et al.* Neutron activation analysis in tobacco and cigarette smoke studies: The halogens. *Beitr Tabakforsch Int* 1982; 11: 195-202.
- 72 Kopplin A *et al.* Urinary excretion of 3-methyladenine and 3-ethyladenine after controlled exposure to tobacco smoke. *Carcinogenesis* 1995; 16: 2637-2641.
- 73 Shigenaga MK, Aboujaoude EN, Chen Q, Ames BN. Assays of oxidative DNA damage biomarkers 8-oxo-2'-deoxyguanosine and 8-oxoguanine in nuclear DNA and biological fluids by high-performance liquid chromatography with electrochemical detection. *Methods Enzymol* 1994; 234: 16-33.
- 74 van Welie RTH, van Dijk RGJM, Vermeulen NPE. Mercapturic acids, protein adducts, and DNA adducts as biomarkers of electrophilic chemicals. *Crit Rev Toxicol* 1992; 22: 271-306.
- 75 van Doorn R *et al.* Detection of human exposure to electrophilic compounds by assay of thioether detoxication products in urine. *Ann Occup Hyg* 1981; 24: 77-92.
- 76 Scherer G *et al.* Urinary mutagenicity and thioethers in nonsmokers: Role of environmental tobacco smoke (ETS) and diet. *Mutat Res* 1996; 368: 195-204.
- 77 Husgafvel-Pursiainen K, Sorsa M, Engström K, Einistö P. Passive smoking at work: biochemical and biological measures of exposure to environmental tobacco smoke. *Int Arch Occup Environ Health* 1985; 59: 337-345.
- 78 Kado NY *et al.* Use of a salmonella micro pre-incubation procedure for studying personal exposure to mutagens in environmental tobacco smoke: Pilot study of urine and airborne mutagenicity from passive smoking. In Sandhu SS, DeMarini DM, Mass MJ, Moore MM, Mumford JL (eds): *Short-term bioassays in the analysis of complex environmental mixtures V*. New York, London, Plenum Press, 1987; pp. 375-390.
- 79 Bartsch H *et al.* Carcinogen hemoglobin adducts, urinary mutagenicity, and metabolic phenotype in active and passive cigarette smokers. *J Natl Cancer Inst* 1990; 82: 1826-1831.
- 80 Sasson IM *et al.* Mutagens in human urine: effects of cigarette smoking and diet. *Mutat Res* 1985; 158: 149-157.
- 81 DeMarini DM. Genotoxicity of tobacco smoke and tobacco smoke condensate. *Mutat Res* 1983; 111: 59-89.
- 82 Sorsa M *et al.* Cytogenetic effects of tobacco smoke exposure among involuntary smokers. *Mutat Res* 1989; 222: 111-116.
- 83 Collman GW *et al.* Effects of α -naphthoflavone on levels of sister chromatid exchanges in lymphocytes from active and passive cigarette smokers: Dose-response relationships. *Cancer Res* 1986; 46: 6452-6455.
- 84 Gorgels WJM *et al.* Passive smoking and sister-chromatid exchanges in lymphocytes. *Mutat Res* 1992; 279: 233-238.
- 85 Pasanen M, Pelkonen O. The expression and environmental regulation of P450 enzymes in human placenta. *Crit Rev Toxicol* 1994; 24: 211-229.
- 86 Manchester DK, Jacoby EH. Sensitivity of human placental monooxygenase activity to maternal smoking. *Clin Pharmacol Ther* 1981; 30: 681-692.
- 87 Huel G *et al.* Aryl hydrocarbon hydroxylase activity in human placenta of passive smokers. *Environ Res* 1988; 50: 173-183.

- 88 Whyatt RM *et al.* CYP1A1 messenger RNA levels in placental tissue as a biomarker of environmental exposure. *Cancer Epidemiol Biomarkers Prev* 1995; **4**: 147-153.
- 89 Remmer H. Passively inhaled tobacco smoke: a challenge to toxicology and preventive medicine. *Arch Toxicol* 1987; **61**: 89-104.
- 90 Langness U. Hydroxyprolinausscheidung und Kollagenstoffwechsel. *Dtsch Med Wochenschr* 1970; **50**: 2530-2535.
- 91 Kasuga H. An introduction to the study of smoking using urinary hydroxyproline, in Kasuga H (ed): *Indoor Air Quality*. Berlin, Springer-Verlag, 1990: pp.37-52.
- 92 Adlkofer F, Scherer G, Heller W-D. Hydroxyproline excretion in urine of smokers and passive smokers. *Prev Med* 1984; **13**: 670-679.
- 93 Verplanke AJW *et al.* Hydroxyproline excretion in schoolchildren and its relationship to measures of indoor air pollution. *Int Arch Occup Environ Health* 1987; **59**: 221-231.
- 94 Adgate JL *et al.* Nitrogen dioxide exposure and urinary excretion of hydroxyproline and desmosine. *Arch Environ Health* 1992; **47**: 376-384.
- 95 US Department of Health and Human Services. *The Health Consequences of Smoking: Cardiovascular Disease: A Report of the Surgeon General*. Rockville, Maryland, Public Health Service, 1983.
- 96 Glantz SA, Parmley WW. Passive smoking and heart disease. *JAMA* 1995; **273**: 1047-1053.
- 97 Taylor AE, Johnson DC, Kazemi H. Environmental tobacco smoke and cardiovascular disease. A position paper from the Council on Cardiopulmonary and Critical Care, American Heart Association. *Circulation* 1992; **86**: 699-702.
- 98 Smith CJ *et al.* Urinary thromboxane and prostacyclin are not altered in nonsmokers living with smokers. *Blood* 1997, submitted.
- 99 Sinzinger H, Kefalides A. Passive smoking severely decreases platelet sensitivity to antiaggregatory prostaglandins. *Lancet* 1982; **2**: 392.
- 100 Burghuber OC *et al.* Platelet sensitivity to prostacyclin in smokers and non-smokers. *Chest* 1986; **89**: 34-38.
- 101 Sinzinger H, Virgolini I. Besitzen Passivraucher ein erhöhtes Thromboserisiko? *Wien Klin Wochenschr* 1989; **101**: 694-698.
- 102 Davis JW, Shelton L, Watanabe IS, Arnold J. Passive smoking affects endothelium and platelets. *Arch Intern Med* 1989; **149**: 386-389.
- 103 Howard G *et al.* Active and passive smoking are associated with increased carotid wall thickness. *Arch Intern Med* 1994; **154**: 1277-1282.
- 104 Diez-Roux AV *et al.* The relationship of active and passive smoking to carotid atherosclerosis 12-14 years later. *Prev Med* 1995; **24**: 48-55.
- 105 Iso H *et al.* Passive smoking and plasma fibrinogen concentrations. *Am J Epidemiol* 1996; **144**: 1151-1154.
- 106 Kawachi I, Colditz GA. Invited commentary: Confounding, measurement error, and publication bias in studies of passive smoking. *Am J Epidemiol* 1996; **144**: 909-915.
- 107 Feldman J *et al.* Passive smoking alters lipid profiles in adolescents. *Pediatrics* 1991; **88**: 259-264.
- 108 Matanoski G, Kanchanaraksa S, Lantry D, Chang Y. Characteristics of nonsmoking women in NHANES I and NHANES I epidemiologic follow-up study with exposure to spouses who smoke. *Am J Epidemiol* 1995; **142**: 149-157.
- 109 Svendsen KH, Kuller LH, Martin MJ, Ockene JK. Effects of passive smoking in the multiple risk factor intervention trial. *Am J Epidemiol* 1987; **126**: 783-795.
- 110 Szadkowski D, Harke HP, Angerer J. Kohlenmonoxydbelastung durch Passivraucher in Büroräumen. *Ann Med* 1976; **3**: 310-313.
- 111 Friedman GD, Petitti DB, Bawol RD. Prevalence and correlates of passive smoking. *Am J Publ Health* 1983; **73**: 401-405.
- 112 Jarvis M *et al.* Biochemical markers of smoke absorption and self reported exposure to passive smoking. *J Epidemiol Commun Health* 1984; **38**: 335-339.
- 113 Tunstall-Pedoe H, Woodward M, Brown CA. Tea drinking, passive smoking, smoking deception and serum cotinine in the Scottish Heart Health Study. *J Clin Epidemiol* 1991; **44**: 1411-1414.
- 114 Scherer G, Conze C, Angerer J, Adlkofer F. Non-occupational exposure to benzene, in Mohr U (ed): *Advances in Controlled Clinical Inhalation Studies*. Berlin, Heidelberg, Springer-Verlag, 1993: pp.387-392.
- 115 Wallace LA *et al.* Exposures to benzene and other volatile compounds from active and passive smoking. *Arch Environ Health* 1987; **42**: 272-279.
- 116 Buratti M *et al.* Evaluation of urinary t,t-muconic acid as a biomarker of benzene exposure related to ETS, in Maroni M (ed): *healthy buildings '95 an international conference on healthy buildings in mild climate*. Milano, healthy buildings '95, 1995: vol 2, pp.605-609.
- 117 MacIure M *et al.* Elevated blood levels of carcinogens in passive smokers. *Am J Publ Health* 1989; **79**: 1381-1384.
- 118 Celermajer DS *et al.* Passive smoking and impaired endothelium-dependent arterial dilatation in healthy young adults. *New Engl J Med* 1996; **334**: 150-154.
- 119 Moskowitz WB *et al.* Lipoprotein and oxygen transport alterations in passive smoking preadolescent children. The MCV Twin Study. *Circulation* 1990; **81**: 586-592.